

Inhibition of protein phosphorylation promotes optic nerve regeneration after injury

Novel approach to the development of treatments for optic neuropathies

Date: May 21, 2019

Source: Waseda University

Summary: Research results suggest that the inhibition of phosphorylation of microtubule-binding protein CRMP2 could be a novel approach to the development of treatments for optic neuropathies, such as glaucoma and traumatic injury.

FULL STORY

A new study by Professor Toshio Ohshima of Waseda University has found that the inhibition of phosphorylation of collapsing response mediator protein 2 (CRMP2), a microtubule-binding protein, suppresses degeneration of nerve fibers and promotes its regeneration after optic nerve injury.

The findings of this study, recently published online in *Scientific Reports*, could be translated into the development of novel treatments for patients with optic neuropathies, such as traumatic injury and glaucoma.

When diagnosed with glaucoma, blind spots develop in the visual field and could lead to blindness as the optic nerve deteriorates. Such deterioration and loss of function of nerve fibers, or axonal degeneration, also occur in traumatic nerve injury and neurological diseases, such as Alzheimer's disease and ALS. Currently, methods for completely repairing the retina, spinal cord, and other parts of the central nervous system after injury or degeneration do not exist because axonal regeneration is restricted by inhibitory factors.

In past studies, potential mechanisms which inhibit axonal regeneration have been found, and addressing these mechanisms were thought to bring scientists a step closer in developing new therapeutics for damages to the central nervous system.

The CRMP2 protein molecule functions to stabilize microtubules, which provide structural support for the central nervous system at the neuronal cellular level and promote polymerization by binding to tubulin dimers. However, these functions are prevented by various kinases through phosphorylation, a mechanism regulating neuronal proteins.

"In our previous study, what we did was to develop CRMP2 knock-in mice and genetically inhibit its CRMP2 phosphorylation," explains Professor Ohshima. "As a result, the CRMP2 knock-in mice showed promotion of axonal regeneration after a spinal cord injury. From this, we hypothesized that the same phenomenon could also be observed after optic nerve injury."

To investigate, the scientists compared the degeneration and regeneration of the optic nerve between wildtype and CRMP2 knock-in mice after an optic nerve injury introduced by an optic nerve crush, a procedure in which the left optic nerve was exposed and crushed for 5 seconds with a tweezer at a site approximately 1mm behind the eye's

globe. They found that destabilization and depolymerization of microtubules after an optic nerve crush injury was suppressed in CRMP2 knock-in mice, and the loss of retinal ganglion cells, the projection neuron of the eye, was also reduced.

The research team also observed that the protein level of GAP43, a molecular marker for axonal regeneration, was higher in the optic nerve from CRMP2 knock-in mice than that from wildtype mice 4 weeks after the optic nerve crush. Additionally, the number of axons increased in the optic nerve after the optic nerve crush in CRMP2 knock-in mice.

"Our experimental data indicate that inhibition of CRMP2 phosphorylation will be a novel approach to the development of treatments for human optic nerve injuries, but further study will be necessary for practical application," says Professor Ohshima.

"For instance, some studies suggest that CRMP2 phosphorylation plays a role in forming myelin, an important substance for sufficient functional recovery of regenerated axons. Additional treatment to facilitate myelination will be required to realize functional recovery after optic nerve injury."

Story Source:

Materials provided by **Waseda University**. *Note: Content may be edited for style and length.*

Journal Reference:

1. Shunsuke Kondo, Kazuya Takahashi, Yuki Kinoshita, Jun Nagai, Shuji Wakatsuki, Toshiyuki Araki, Yoshio Goshima, Toshio Ohshima. **Genetic inhibition of CRMP2 phosphorylation at serine 522 promotes axonal regeneration after optic nerve injury**. *Scientific Reports*, 2019; 9 (1) DOI: 10.1038/s41598-019-43658-w

Cite This Page:

[MLA](#)[APA](#)[Chicago](#)

Waseda University. "Inhibition of protein phosphorylation promotes optic nerve regeneration after injury: Novel approach to the development of treatments for optic neuropathies." *ScienceDaily*. ScienceDaily, 21 May 2019. <www.sciencedaily.com/releases/2019/05/190521101453.htm>.

RELATED STORIES

Every Cell Has a Story to Tell in Brain Injury

Oct. 4, 2018 — Traumatic head injury can have widespread effects in the brain, but now scientists can look in real time at how head injury affects thousands of individual cells and genes simultaneously in mice. ... **read more »**

Timing Is Key for Traumatic Brain Injury Treatment

Oct. 14, 2014 — Two potential treatments for traumatic brain injury have been found to be most effective when given at different stages after the injury has occurred. Laboratory studies have confirmed that changes ... **read more »**

Second Look at Glaucoma Surgery: Anti-Inflammatory Medications After Glaucoma Laser Surgery Not Helpful or Necessary

Sep. 18, 2014 — Using anti-inflammatory medications after glaucoma laser surgery is not helpful or necessary, new research concludes. Glaucoma is the most common cause of irreversible blindness in the world, and is ...

read more »

Potential New Therapy With Brain-on-a-Chip Axonal Strain Injury Model

July 17, 2014 — The use of 'Brain-on-a-Chip' microsystem has been used to assess specific effects of traumatic axonal injury. This innovative approach was used to characterize the biochemical changes that are ... **read more »**